arXiv:1212.5037v1 [hep-ph] 20 Dec 2012

Fragmentation contribution to the transverse single-spin asymmetry in proton-proton collisions

A. Metz and D. Pitonyak

Department of Physics, Barton Hall, Temple University, Philadelphia, PA 19122, USA

December 21, 2012

Abstract

Within the collinear twist-3 framework, we study the single-spin asymmetry (SSA) in collisions between unpolarized protons and transversely polarized protons with focus on the fragmentation term. The fragmentation mechanism must be analyzed in detail in order to unambiguously determine the impact of various contributions to SSAs in hadron production. Such a distinction may also settle the "sign mismatch" between the transverse SSA in proton-proton collisions and the Sivers effect in semi-inclusive deep inelastic scattering. We calculate terms involving quark-quark and quark-gluon-quark correlators, which is an important step in such an investigation.

1 Introduction

Beginning in the late 1970s, calculations of transverse SSAs in inclusive hadron production within the naïve collinear parton model demonstrated that such asymmetries should be on the order of $\alpha_s m_q/P_{h\perp}$, where m_q is the mass of the quark and $P_{h\perp}$ is the transverse momentum of the detected hadron [1,2]. These predictions contradict the large SSAs seen in experiments [3–9]. However, the use of collinear twist-3 multi-parton correlators established a framework, valid when $\Lambda_{QCD} \ll P_{h\perp}$, that could potentially handle these observables [10–12]. Various processes have been analyzed over the last two decades using this methodology — see [11–23] for some specific examples. (We also mention that other mechanisms have been proposed to explain large SSAs [24–27].) Furthermore, the collinear twist-3 formalism has been used to describe the double-spin observable A_{LT} in different reactions with one large scale [28–35].

In particular, much attention has been given to the transverse target SSA for inclusive single hadron production in proton-proton collisions. This observable was initially believed to be dominated by soft-gluon-pole (SGP) contributions on the side of the transversely polarized nucleon [12,14], which involves the Efremov-Teryaev-Qiu-Sterman (ETQS) function $T_F(x,x)$. However, a recent fit of $T_F(x,x)$ to $pp^\uparrow \to \pi X$ SSA data based on this assumption [14] has led to the so-called "sign mismatch" crisis [36] involving the ETQS function and the transverse momentum dependent (TMD) Sivers function extracted from semi-inclusive deep inelastic scattering (SIDIS) [37] — see also the discussion in [21]. A conceivable resolution could be that soft-fermion-poles and/or tri-gluon correlations in the nucleon provide a significant effect [16, 19]. One also cannot rule out the importance of the fragmentation mechanism, which could at the very least give a piece comparable to SGP contributions — see [38–40] and references therein. These additional terms may be able to account for the sign mismatch. Other possible explanations have also been explored [36, 41].

Actually, the fragmentation contribution has a counterpart in the TMD factorization approach known as the Collins mechanism [42]. This causes azimuthal modulations in the cross section for processes like SIDIS and $e^+e^- \to h_1 h_2 X$. These effects have been measured in both processes [43–46], which has allowed for an extraction of the Collins function [47–49]. In addition, both the collinear twist-3 and TMD approaches to SSAs from fragmentation in SIDIS have been shown to agree in an intermediate momentum region where both formalisms are valid [50]. The TMD Collins mechanism has also been used to describe the fragmentation piece of the SSA in $pp^\uparrow \to \pi X$ [39, 40], although a rigorous proof has not been put forth that such a framework can be applied to a process with one large scale.

Within the collinear twist-3 approach, attempts have been made to formulate the fragmentation term in the SSA for inclusive pion production from proton-proton collisions [51]. However, further work determined the contribution considered in Ref. [51] vanishes due to a universality argument [52–56]. Recent calculations have rectified the situation, and the so-called derivative term was computed for the first time in Ref. [38]. In the current work, we derive not only this term but also the non-derivative term as well as contributions involving quark-gluon-quark (qgq) correlators. In addition, we briefly comment on the implications of this and future studies on the resolution of the "sign mismatch" puzzle as well as the overall understanding of SSAs in proton-proton collisions.

The paper is organized as follows: in Sec. 2 we review the collinear twist-3 formalism, and, in particular, discuss the relevant unpolarized fragmentation functions (FFs). In Sec. 3 we present the result for the fragmentation contribution to the single-spin dependent cross section in $pp^\uparrow \to hX$ and give a few details of the calculation. We conclude the paper in Sec. 4.

2 Collinear twist-3 formalism and unpolarized FFs

To start, let us make explicit the process under consideration, namely,

$$A(P, \vec{S}_\perp) + B(P') \to C(P_h) + X, \tag{1}$$

where the 4-momenta and polarizations of the incoming protons A, B and outgoing hadron C are indicated. The Mandelstam variables for the process are defined as $S = (P + P')^2$, $T = (P - P_h)^2$, and $U = (P' - P_h)^2$, which on the partonic level give $\hat{s} = xx'S$, $\hat{t} = xT/z$, and $\hat{u} = x'U/z$. The longitudinal momentum fraction x (x') is associated with partons in the transversely polarized (unpolarized) nucleon. In analogy to the usually defined lightcone vectors $n = (0^+, 1^-, 0_\perp)$ and $\bar{n} = (1^+, 0^-, 0_\perp)$, we also have $n_h \sim P_h$ and $\bar{n}_h \sim (P_h^0, -\vec{P}_h)$ (with $n_h \cdot \bar{n}_h = 1$) as lightcone vectors associated with the outgoing hadron's direction of motion [38]. Such vectors allow us to perform the appropriate twist expansion of the fragmentation correlator in the context of this process. We also note that $\epsilon_\perp^{\mu\nu} = \epsilon^{\rho\sigma\mu\nu}\bar{n}_\rho n_\sigma$ with $\epsilon_\perp^{12} = 1$. We perform the calculation of the transverse SSA in the proton-proton *cm*-frame, with the transversely polarized proton moving along the positive z-axis.

The first non-vanishing contribution to the spin-dependent cross section is given by terms of twist-3 accuracy and reads

$$\begin{aligned} d\sigma(\vec{P}_{h\perp}, \vec{S}_\perp) = \; & H \otimes f_{a/A(3)} \otimes f_{b/B(2)} \otimes D_{C/c(2)} \\ & + H' \otimes f_{a/A(2)} \otimes f_{b/B(3)} \otimes D_{C/c(2)} \\ & + H'' \otimes f_{a/A(2)} \otimes f_{b/B(2)} \otimes D_{C/c(3)}, \end{aligned} \tag{2}$$

where a sum over partonic channels and parton flavors in each channel is understood. In Eq. (2), $f_{a/A(t)}$ ($f_{b/B(t)}$) denotes the distribution function associated with parton a (b) in hadron A (B), while $D_{C/c(t)}$ represents the fragmentation function associated with particle C in parton c. The twist of the functions is indicated by t. The factors H, H', and H'' give the hard parts corresponding to each term, while the symbol $\otimes$ denotes convolutions in the appropriate momentum fractions. As



Figure 1: Feynman diagrams for the twist-3 matrix elements that give contributions to $D_{C/c(3)}$. See the text for more details.

mentioned, the first term in (2) has been analyzed previously in the literature [12, 14–16]. Likewise, the second term, which involves chiral-odd twist-3 unpolarized distributions, has also been studied and was shown to be negligible because of the smallness of the hard scattering coefficients [57]. The focus of this work will be on the third term in Eq. (2) involving collinear twist-3 FFs. Therefore, for the situation we consider, $f_{a/A(2)} = h_1^a$ and $f_{b/B(2)} = f_1^b$, where h_1 and f_1 are the standard twist-2 transversity distribution function and unpolarized distribution function, respectively. We then must determine what contributions are possible for $D_{C/c(3)}$.

For a detailed discussion of collinear twist-3 distribution functions see, for instance, Ref. [18]; the same formalism can be generalized to the fragmentation case. Here we consider the situation where the outgoing hadron has a large "minus" component of momentum. The twist-3 matrix elements that we must consider are given by the diagrams in Fig. 1. In the lightcone ($A^- = 0$) gauge, these graphs lead to the three matrix elements

$$\langle \psi | \, \rangle \langle \, | \bar{\psi} \rangle, \; \langle \partial_\perp \psi | \, \rangle \langle \, | \bar{\psi} \rangle, \; \langle A_\perp \psi | \, \rangle \langle \, | \bar{\psi} \rangle, \tag{3}$$

which result from Figs. 1(a), (b), and (d), respectively. The symbol $| \, \rangle \langle \, |$ represents the intermediate $|P_h; X\rangle \langle P_h; X|$ in the fragmentation correlators. We do not have to consider Fig. 1(c) because one does not need to simultaneously take into account $k_\perp$ expansion and $A_\perp$ gluon attachments (which would give rise to twist-4 contributions).

Now that we have determined the relevant twist-3 matrix elements, we must parameterize them in terms of twist-3 FFs that will eventually be involved in our final result. We will follow the work of Ref. [58] in defining these FFs and the relations between them. We first focus on the qgq matrix element $\langle A_\perp \psi | \, \rangle \langle \, | \bar{\psi} \rangle$. One notices that this matrix element is not gauge invariant. In analogy to the qgq distribution correlators, this can be resolved in two ways: rewrite the gluon field $A_\perp$ in terms of the field strength tensor $F_\perp^{-\mu} = \partial^- A_\perp^\mu$ or rewrite it in terms of the covariant derivative $D_\perp^\mu = \partial_\perp^\mu - igA_\perp^\mu$. Thus, we also have so-called "F-type" and "D-type" functions on the fragmentation side. For unpolarized final-state hadrons, these read

$$\sum_X\!\!\!\!\!\!\!\int \frac{1}{z}\int\frac{d\xi^+}{2\pi}\int\frac{d\zeta^+}{2\pi}\,e^{i\frac{P_h^-}{z_1}\xi^+}\,e^{i(\frac{1}{z}-\frac{1}{z_1})P_h^-\zeta^+}\langle 0|igF_\perp^{-\mu}(\zeta^+)\psi_\alpha(\xi^+)|P_h;X\rangle\langle P_h;X|\bar\psi_\beta(0)|0\rangle$$
$$= M_h\left[\epsilon_\perp^{\mu\nu}\,\sigma_\nu{}^+\gamma_5\,\hat H^q_{FU}(z,z_1)\right]_{\alpha\beta}, \tag{4}$$

and

$$\sum_X\!\!\!\!\!\!\!\int \frac{1}{z}\int\frac{d\xi^+}{2\pi}\int\frac{d\zeta^+}{2\pi}\,e^{i\frac{P_h^-}{z_1}\xi^+}\,e^{i(\frac{1}{z}-\frac{1}{z_1})P_h^-\zeta^+}\langle 0|iD_\perp^{\mu}(\zeta^+)\psi_\alpha(\xi^+)|P_h;X\rangle\langle P_h;X|\bar\psi_\beta(0)|0\rangle$$
$$= \frac{M_h}{P_h^-}\left[\epsilon_\perp^{\mu\nu}\,\sigma_\nu{}^+\gamma_5\,\hat H^q_{DU}(z,z_1)\right]_{\alpha\beta}. \tag{5}$$

We remark that these functions contain both real and imaginary parts, which we respectively indicate by $\hat H^{q,\mathfrak{R}}_{FU(DU)}(z,z_1)$ and $\hat H^{q,\mathfrak{I}}_{FU(DU)}(z,z_1)$. In Eqs. (4), (5), we have suppressed Wilson lines and have indicated the hadron mass by M_h. Moreover, the F-type and D-type functions are not independent of each other. One can establish the following relation between them:

$$\hat H^q_{DU}(z,z_1) = -\frac{i}{z^2}\hat H^q(z)\,\delta\left(\frac{1}{z}-\frac{1}{z_1}\right) + PV\frac{1}{\frac{1}{z}-\frac{1}{z_1}}\hat H^q_{FU}(z,z_1), \tag{6}$$

where PV denotes the principal value. In order to derive this expression, notice that we must introduce an additional twist-3 function $\hat H^q(z)$, which is given by

$$\sum_X\!\!\!\!\!\!\!\int z\int\frac{d\xi^+}{2\pi}e^{i\frac{P_h^-}{z}\xi^+}\langle 0|\left(iD_\perp^\mu(\xi^+)+g\int_{\xi^+}^{\infty}d\zeta^+F_\perp^{-\mu}(\zeta^+)\right)\psi_\alpha(\xi^+)|P_h;X\rangle\langle P_h;X|\bar\psi_\beta(0)|0\rangle$$
$$= -iM_h\left[\epsilon_\perp^{\mu\nu}\,\sigma_\nu{}^+\gamma_5\,\hat H^q(z)\right]_{\alpha\beta}. \tag{7}$$

This function is associated with the quark-quark (qq) matrix element $\langle\partial_\perp\psi|\,\rangle\langle\,|\bar\psi\rangle$. We also mention that $\hat H(z)$ is equivalent to the first $k_\perp$-moment of the Collins function $H_1^\perp(z,z^2\vec k_\perp^2)$ (as defined in [59, 60]) for the fragmentation of a transversely polarized quark into an unpolarized hadron:

$$\hat H^q(z) = z^2\int d^2\vec k_\perp\,\frac{\vec k_\perp^{\,2}}{2M_h^2}\,H_1^{\perp,q}(z,z^2\vec k_\perp^2). \tag{8}$$

We note that $\hat H^q(z)$ is a real valued function, so that Eq. (6) in particular tells us

$$\hat H^{q,\mathfrak{I}}_{DU}(z,z_1) = -\frac{1}{z^2}\hat H^q(z)\,\delta\left(\frac{1}{z}-\frac{1}{z_1}\right) + PV\frac{1}{\frac{1}{z}-\frac{1}{z_1}}\hat H^{q,\mathfrak{I}}_{FU}(z,z_1), \tag{9}$$

$$\hat H^{q,\mathfrak{R}}_{DU}(z,z_1) = PV\frac{1}{\frac{1}{z}-\frac{1}{z_1}}\hat H^{q,\mathfrak{R}}_{FU}(z,z_1). \tag{10}$$

The other relevant qq matrix element $\langle\psi|\,\rangle\langle\,|\bar\psi\rangle$ leads to a contribution from the twist-3 function $H^q(z)$, whose definition is given by [59]

$$\sum_X\!\!\!\!\!\!\!\int z\int\frac{d\xi^+}{2\pi}e^{i\frac{P_h^-}{z}\xi^+}\langle 0|\psi_\alpha(\xi^+)|P_h;X\rangle\langle P_h;X|\bar\psi_\beta(0)|0\rangle = -\frac{iM_h}{2P_h^-}\left[\epsilon_\perp^{\mu\nu}\,\sigma_{\mu\nu}\gamma_5\,H^q(z)\right]_{\alpha\beta}. \tag{11}$$



Figure 2: Graphs showing factorization for fragmentation contributions to A_{UT} from (a) qq correlators and (b) qgq correlators.

However, $H^q(z)$ can be related to the imaginary part of the D-type function through the QCD equations of motion (EOM):

$$H^q(z) = 2z^3 \int \frac{dz_1}{z_1^2} \hat{H}_{DU}^{q,\Im}(z, z_1). \tag{12}$$

Thus, we see from Eqs. (6), (12) that of the four collinear twist-3 FFs relevant for an unpolarized hadron ($\hat{H}, H, \hat{H}_{FU}, \hat{H}_{DU}$), only two are independent of each other. However, we will find it convenient to write the final result in terms of $\hat{H}$, H, and $\hat{H}_{FU}$. In fact, it turns out only $\hat{H}_{FU}^{\Im}$ is relevant for the calculation, as the contributions to $\hat{H}_{FU}^{\Re}$ vanish.

3 Calculation of the single spin-dependent cross section

The derivation of the fragmentation contribution to the SSA in $pp^\uparrow \to hX$ follows a similar procedure to what was outlined in Ref. [35] for the double-spin asymmetry A_{LT} in the same process. The factorization of the reaction under consideration is shown in Fig. 2. This includes collinear factors associated with the upolarized nucleon (top gray blob), the outgoing hadron (middle gray blob), and the transversely polarized nucleon (bottom gray blob) as well as hard factors (white blobs). For each partonic channel, the main task becomes calculating the hard scattering coefficients for each of these functions, which then allows us to write down the single-spin dependent cross section. We will denote each channel by $ab \to cd$, where $a\,(b)$ is the parton associated with the transversely polarized (unpolarized) nucleon and c is the parton that fragments into the detected hadron.

If we keep the transverse momentum (relative to the outgoing hadron) of the parton c (as in Fig. 1(b)), then we can determine the hard scattering coefficient for $\hat{H}(z)$ and its derivative. On the other hand, if we neglect the transverse momentum of c in the initial state (as in Fig. 1(a)), then we obtain the hard part for $H(z)$. Lastly, we must attach gluons to the hard blobs, in which case we can neglect the transverse momentum of the parton c and gluon (as in Fig. 1(d)). These diagrams allow us to find the qgq contributions to the hard factors for $\hat{H}_{FU}^{\Im}$. We remark at this stage that in general the qgq diagrams are not always real but can acquire an imaginary part whenever internal parton lines go on-shell. This requires the use of the distribution identity

$$\frac{1}{x \pm i\epsilon} = PV\frac{1}{x} \mp i\pi\delta(x). \tag{13}$$

As was the case in [34, 35], only the PV part survives the sum over the various cut diagrams. These qgq hard parts contain both z_1-dependent and -independent terms. The latter can be pulled out of the integral over z_1, which allows $\hat{H}_{FU}^{\Im}$ for those pieces to be rewritten in terms of $\hat{H}$ and H. In the end, we find the fragmentation contribution to the cross section relevant for the SSA A_{UT} in $AB \to CX$ is given by

$$\frac{P_h^0\, d\sigma(\vec{S}_\perp)}{d^3\vec{P}_h} = -\frac{2\alpha_s^2 M_h}{S}\epsilon_{\perp,\alpha\beta}\, S_\perp^\alpha P_{h\perp}^\beta \sum_i \sum_{a,b,c} \int_{z_{min}}^1 \frac{dz}{z^3} \int_{x'_{min}}^1 \frac{dx'}{x'}\,\frac{1}{x}\,\frac{1}{x'S+T/z}\,\frac{1}{-x'\hat{t}-x\hat{u}}\, h_1^a(x)\, f_1^b(x')$$
$$\times \left\{ \left[\hat{H}^c(z) - z\frac{d\hat{H}^c(z)}{dz}\right] S_{\hat{H}}^i + \frac{1}{z} H^c(z)\, S_H^i \right.$$
$$\left. + 2z^2 \int \frac{dz_1}{z_1^2} PV \frac{1}{\frac{1}{z}-\frac{1}{z_1}} \hat{H}_{FU}^{c,\Im}(z,z_1)\, \frac{1}{\xi}\, S_{\hat{H}_{FU}}^i \right\}, \qquad (14)$$

where i denotes the channel, $x = -x'(U/z)/(x'S+T/z)$, $x'_{min} = -(T/z)/(U/z+S)$, and $z_{min} = -(T+U)/S$. We have introduced $\xi = z/z_g$, where $z_g = zz_1/(z_1 - z)$, and understand $1/\xi$ to mean $PV(1/\xi)$. The hard scattering coefficients S^i are given in the Appendix.

A few comments are in order on the final result. First, note that the number of possible channels reduces significantly due to vanishing Dirac traces that arise in certain partonic interactions. Second, we remark that the "compact" form involving $\hat{H}$ and its derivative manifest in Eq. (14) has been achieved by writing the cross section in terms of $\hat{H}$, H, and $\hat{H}_{FU}^{\Im}$. Such a structure has arisen in other spin-dependent cross sections [11, 14, 15, 32–35]. In particular, we point out that unlike Refs. [34, 35], the compact form in (14) was obtained by writing the result using F-type functions rather than D-type functions. We repeat that the derivative term was first computed in Ref. [38].

4 Summary and outlook

In conclusion, we have calculated the fragmentation contribution to the SSA in hadron production from proton-proton collisions using collinear twist-3 factorization, including terms involving quark-quark and quark-gluon-quark correlators. The derivative term in this context had been computed before [38]. We generalized the work in Ref. [18] to the fragmentation case and specified the relevant collinear twist-3 functions — see also [38, 50]. The structure of the calculation essentially follows the procedure outlined in [35] for the double-spin asymmetry A_{LT} in the same process. Moreover, we find a "compact" form for $\hat{H}$ and its derivative that has also arisen in other spin-dependent cross sections [11, 14, 15, 32–35].

The fragmentation mechanism is a key piece to this observable, and an estimate of its impact on the asymmetry had been considered previously [38–40]. However, in order to completely determine the size of this term using the collinear twist-3 formalism, a full study including quark-gluon-quark correlators will be necessary. The current work is an important step in this investigation. In the end, a global analysis involving several processes will be required in order to extract the twist-3 functions (both distribution and fragmentation) that enter into the SSA in $pp^\uparrow \to hX$. Nevertheless, given the recent "sign mismatch" crisis involving the ETQS function and the Sivers function [36], such an intensive study is worthwhile in order to thoroughly understand the mechanism behind this observable and whether the collinear twist-3 framework can appropriately describe it.

Acknowledgments: We would like to thank Z.-B. Kang and J. Zhou for useful discussions with regards to Ref. [38]. This work has been supported by the NSF under Grant No. PHY-1205942.

Appendix: Hard scattering coefficients

Here we give the hard scattering coefficients S^i found in Eq. (14). We mention that the $SU(3)$ color factors depend on $N_c = 3$.

$qg \to qg$ channel

$$S_{\hat{H}} = (x - x')\left[\frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} - \frac{1}{N_c^2}\right] \tag{15}$$

$$S_H = -\frac{1}{2}\left[\frac{\hat{t}^3(x'\hat{t} - x\hat{u}) - 6x'\hat{t}^2\hat{u}(\hat{s} - \hat{u}) + 4\hat{u}^3(2x'\hat{t} - x\hat{s})}{\hat{t}^3\hat{u}}\right] + \frac{1}{N_c^2}\left[\frac{x'\hat{s} + (x - x')\hat{u}}{\hat{u}}\right] \tag{16}$$

$$S_{\hat{H}_{FU}} = \frac{1}{N_c^2 - 1}\left[\frac{\hat{s}(\hat{s} - \hat{u})(-x'\hat{t} - x\hat{u})}{\hat{t}^3}\right] + \frac{1}{N_c^2}\left[\frac{x\hat{s}}{\hat{t}}\right] + \frac{x'\hat{s}^3 + (x - x')\hat{u}^3}{\hat{t}^2\hat{u}} \tag{17}$$

$qq \to qq$ channel

$$S_{\hat{H}} = \frac{N_c^2 - 1}{N_c^2}(x - x')\left\{-\frac{\hat{s}\hat{u}}{\hat{t}^2} + \frac{1}{N_c}\left[\frac{\hat{s}}{\hat{t}}\right]\right\} \tag{18}$$

$$S_H = \frac{N_c^2 - 1}{N_c^2}\left\{\frac{\hat{s}(x'\hat{s}^2 + (x - x')\hat{u}^2)}{\hat{t}^3} - \frac{1}{N_c}\left[\frac{\hat{s}(x'\hat{s}^2 - (x - x')\hat{u}(\hat{t} - \hat{u}))}{2\hat{t}^2\hat{u}}\right]\right\} \tag{19}$$

$$S_{\hat{H}_{FU}} = -\frac{x'\hat{s}^2}{\hat{t}^2} + \frac{1}{N_c}\left[\frac{(x - x')\hat{s}}{\hat{t}}\right] - \frac{1}{N_c^2}\left[\frac{\hat{s}(2x'\hat{t}^2 + x\hat{u}(\hat{t} - \hat{u}))}{\hat{t}^3}\right] - \frac{1}{N_c^3}\left[\frac{\hat{s}(x\hat{u}^2 - x'\hat{t}^2)}{\hat{t}^2\hat{u}}\right] \tag{20}$$

$q\bar{q} \to q\bar{q}$ channel

$$S_{\hat{H}} = \frac{N_c^2 - 1}{N_c^2}(x - x')\left\{-\frac{\hat{s}\hat{u}}{\hat{t}^2} + \frac{1}{N_c}\left[\frac{\hat{u}}{\hat{t}}\right]\right\} \tag{21}$$

$$S_H = \frac{N_c^2 - 1}{N_c^2}\left\{\frac{\hat{s}(x'\hat{s}^2 + (x - x')\hat{u}^2)}{\hat{t}^3} - \frac{1}{N_c}\left[\frac{x'\hat{s}(\hat{s} - \hat{t}) + (x - x')\hat{u}^2}{2\hat{t}^2}\right]\right\} \tag{22}$$

$$S_{\hat{H}_{FU}} = -\frac{(x - x')\hat{s}\hat{u}}{\hat{t}^2} + \frac{1}{N_c}\left[\frac{x'\hat{s}}{\hat{t}}\right] + \frac{1}{N_c^2}\left[\frac{\hat{s}(2x'\hat{t}^2 - x\hat{u}(\hat{s} - 2\hat{t}))}{\hat{t}^3}\right] - \frac{1}{N_c^3}\left[\frac{x\hat{s}^2 - (x - x')\hat{t}^2}{\hat{t}^2}\right] \tag{23}$$

$\bar{q}q \to q\bar{q}$ channel

$$S_{\hat{H}} = -\frac{N_c^2 - 1}{N_c^3}(x - x') \tag{24}$$

$$S_H = -\frac{N_c^2 - 1}{N_c^3}\left[\frac{x'\hat{s} + (x - x')\hat{u}}{2\hat{u}}\right] \tag{25}$$

$$S_{\hat{H}_{FU}} = \frac{1}{N_c}\left[\frac{x\hat{s}}{\hat{t}}\right] - \frac{1}{N_c^3}\left[\frac{x'\hat{s}^2 + (x - x')\hat{u}^2}{\hat{t}\hat{u}}\right] \tag{26}$$

$\underline{qq' \to qq'}$ channel

$$S_{\hat{H}} = \frac{N_c^2-1}{N_c^2}(x-x')\left[-\frac{\hat{s}\hat{u}}{\hat{t}^2}\right] \tag{27}$$

$$S_H = \frac{N_c^2-1}{N_c^2}\left[\frac{\hat{s}(x'\hat{s}^2+(x-x')\hat{u}^2)}{\hat{t}^3}\right] \tag{28}$$

$$S_{\hat{H}_{FU}} = -\frac{x'\hat{s}^2}{\hat{t}^2} - \frac{1}{N_c^2}\left[\frac{\hat{s}(2x'\hat{t}^2+x\hat{u}(\hat{t}-\hat{u}))}{\hat{t}^3}\right] \tag{29}$$

$\underline{q\bar{q}' \to q\bar{q}'}$ channel

$$S_{\hat{H}} = S_{\hat{H}}^{qq'\to qq'} \tag{30}$$

$$S_H = S_H^{qq'\to qq'} \tag{31}$$

$$S_{\hat{H}_{FU}} = -\frac{(x-x')\hat{s}\hat{u}}{\hat{t}^2} + \frac{1}{N_c^2}\left[\frac{\hat{s}(2x'\hat{t}^2-x\hat{u}(\hat{s}-2\hat{t}))}{\hat{t}^3}\right] \tag{32}$$

Note that we can obtain the channels involving the fragmentation of an antiquark by charge conjugating the above partonic processes $ab \to cd$. The respective hard parts are given by

$\underline{\bar{a}\bar{b} \to \bar{c}\bar{d}}$ channels

$$S_{\hat{H}} = S_{\hat{H}}^{ab\to cd} \tag{33}$$

$$S_H = S_H^{ab\to cd} \tag{34}$$

$$S_{\hat{H}_{FU}} = -S_{\hat{H}_{FU}}^{ab\to cd} \tag{35}$$

References

[1] G. L. Kane, J. Pumplin and W. Repko, Phys. Rev. Lett. **41**, 1689 (1978).

[2] J. P. Ma and H. Z. Sang, JHEP **0811**, 090 (2008) [arXiv:0809.4811 [hep-ph]]; Phys. Lett. B **676**, 74 (2009) [arXiv:0811.0224 [hep-ph]].

[3] G. Bunce *et al.*, Phys. Rev. Lett. **36**, 1113 (1976).

[4] D. L. Adams *et al.* [E581 and E704 Collaborations], Phys. Lett. B **261**, 201 (1991); D. L. Adams *et al.* [E704 Collaboration], Phys. Lett. B **264**, 462 (1991).

[5] K. Krueger *et al.*, Phys. Lett. B **459**, 412 (1999).

[6] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 171801 (2004) [hep-ex/0310058]; B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **101**, 222001 (2008) [arXiv:0801.2990 [hep-ex]].

[7] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **95**, 202001 (2005) [hep-ex/0507073].

[8] I. Arsene *et al.* [BRAHMS Collaboration], Phys. Rev. Lett. **101**, 042001 (2008) [arXiv:0801.1078 [nucl-ex]].

[9] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. D **86**, 051101 (2012) [arXiv:1205.6826 [nucl-ex]].

[10] A. V. Efremov and O. V. Teryaev, Sov. J. Nucl. Phys. **36**, 140 (1982) [Yad. Fiz. **36**, 242 (1982)]; Phys. Lett. B **150**, 383 (1985).

[11] J.-w. Qiu and G. F. Sterman, Phys. Rev. Lett. **67**, 2264 (1991); Nucl. Phys. B **378**, 52 (1992).

[12] J.-w. Qiu and G. F. Sterman, Phys. Rev. D **59**, 014004 (1999) [hep-ph/9806356].

[13] H. Eguchi, Y. Koike and K. Tanaka, Nucl. Phys. B **752**, 1 (2006) [arXiv:hep-ph/0604003]; Nucl. Phys. B **763**, 198 (2007) [arXiv:hep-ph/0610314].

[14] C. Kouvaris, J. W. Qiu, W. Vogelsang and F. Yuan, Phys. Rev. D **74**, 114013 (2006) [arXiv:hep-ph/0609238].

[15] Y. Koike and K. Tanaka, Phys. Rev. D **76**, 011502 (2007) [hep-ph/0703169].

[16] Y. Koike and T. Tomita, Phys. Lett. B **675**, 181 (2009) [arXiv:0903.1923 [hep-ph]].

[17] W. Vogelsang and F. Yuan, Phys. Rev. D **79**, 094010 (2009) [arXiv:0904.0410 [hep-ph]].

[18] J. Zhou, F. Yuan and Z. T. Liang, Phys. Rev. D **81**, 054008 (2010) [arXiv:0909.2238 [hep-ph]].

[19] K. Kanazawa and Y. Koike, Phys. Rev. D **82**, 034009 (2010) [arXiv:1005.1468 [hep-ph]].

[20] L. Gamberg and Z. -B. Kang, Phys. Lett. B **718**, 181 (2012) [arXiv:1208.1962 [hep-ph]].

[21] A. Metz, D. Pitonyak, A. Schaefer, M. Schlegel, W. Vogelsang and J. Zhou, arXiv:1209.3138 [hep-ph].

[22] M. Schlegel, arXiv:1211.3579 [hep-ph].

[23] Z. -B. Kang, I. Vitev and H. Xing, arXiv:1212.1221 [hep-ph].

[24] P. Hoyer and M. Jarvinen, JHEP **0702**, 039 (2007) [hep-ph/0611293]; P. Hoyer, M. Jarvinen and S. Kurki, JHEP **0810**, 086 (2008) [arXiv:0808.0626 [hep-ph]].

[25] Y. Qian and I. Zahed, Phys. Rev. D **86**, 014033 (2012) [Erratum-ibid. D **86**, 059902 (2012)] [arXiv:1112.4552 [hep-ph]].

[26] Y. V. Kovchegov and M. D. Sievert, Phys. Rev. D **86**, 034028 (2012) [arXiv:1201.5890 [hep-ph]].

[27] S. M. Troshin and N. E. Tyurin, arXiv:1211.4341 [hep-ph].

[28] R. L. Jaffe and X. D. Ji, Phys. Rev. Lett. **67**, 552 (1991); Nucl. Phys. B **375**, 527 (1992).

[29] R. D. Tangerman and P. J. Mulders, hep-ph/9408305.

[30] Y. Koike, K. Tanaka and S. Yoshida, Phys. Lett. B **668**, 286 (2008) [arXiv:0805.2289 [hep-ph]].

[31] Z. Lu and I. Schmidt, Phys. Rev. D **84**, 114004 (2011) [arXiv:1109.3232 [hep-ph]].

[32] A. Metz and J. Zhou, Phys. Lett. B **700**, 11 (2011) [arXiv:1006.3097 [hep-ph]].

[33] Z. B. Kang, A. Metz, J. W. Qiu and J. Zhou, Phys. Rev. D **84**, 034046 (2011) [arXiv:1106.3514 [hep-ph]].

[34] Z. -T. Liang, A. Metz, D. Pitonyak, A. Schafer, Y. -K. Song and J. Zhou, Phys. Lett. B **712**, 235 (2012) [arXiv:1203.3956 [hep-ph]].

[35] A. Metz, D. Pitonyak, A. Schaefer and J. Zhou, arXiv:1210.6555 [hep-ph].

[36] Z. -B. Kang, J. -W. Qiu, W. Vogelsang and F. Yuan, Phys. Rev. D **83**, 094001 (2011) [arXiv:1103.1591 [hep-ph]].

[37] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, S. Melis, F. Murgia, A. Prokudin and C. Turk, Eur. Phys. J. A **39**, 89 (2009) [arXiv:0805.2677 [hep-ph]].

[38] Z. -B. Kang, F. Yuan and J. Zhou, Phys. Lett. B **691**, 243 (2010) [arXiv:1002.0399 [hep-ph]].

[39] Z. -B. Kang and F. Yuan, Phys. Rev. D **84**, 034019 (2011) [arXiv:1106.1375 [hep-ph]].

[40] M. Anselmino, M. Boglione, U. D'Alesio, E. Leader, S. Melis, F. Murgia and A. Prokudin, Phys. Rev. D **86**, 074032 (2012) [arXiv:1207.6529 [hep-ph]].

[41] Z. -B. Kang and A. Prokudin, Phys. Rev. D **85**, 074008 (2012) [arXiv:1201.5427 [hep-ph]].

[42] J. C. Collins, Nucl. Phys. B **396**, 161 (1993) [hep-ph/9208213].

[43] A. Airapetian *et al.* [HERMES Collaboration], Phys. Rev. Lett. **94**, 012002 (2005) [hep-ex/0408013]; Phys. Lett. B **693**, 11 (2010) [arXiv:1006.4221 [hep-ex]]; arXiv:1204.4161 [hep-ex].

[44] K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **96**, 232002 (2006) [hep-ex/0507063]; R. Seidl *et al.* [Belle Collaboration], Phys. Rev. D **78**, 032011 (2008) [Erratum-ibid. D **86**, 039905 (2012)] [arXiv:0805.2975 [hep-ex]].

[45] V. Y. Alexakhin *et al.* [COMPASS Collaboration], Phys. Rev. Lett. **94** (2005) 202002 [hep-ex/0503002]; E. S. Ageev *et al.* [COMPASS Collaboration], Nucl. Phys. B **765**, 31 (2007) [hep-ex/0610068]; M. Alekseev *et al.* [COMPASS Collaboration], Phys. Lett. B **673**, 127 (2009) [arXiv:0802.2160 [hep-ex]]; Phys. Lett. B **692**, 240 (2010) [arXiv:1005.5609 [hep-ex]]; F. Bradamante [COMPASS Collaboration], Nuovo Cim. C **035N2**, 107 (2012) [arXiv:1111.0869 [hep-ex]].

[46] I. Garzia [BABAR Collaboration], arXiv:1211.5293 [hep-ex].

[47] W. Vogelsang and F. Yuan, Phys. Rev. D **72**, 054028 (2005) [hep-ph/0507266].

[48] A. V. Efremov, K. Goeke and P. Schweitzer, Phys. Rev. D **73**, 094025 (2006) [hep-ph/0603054].

[49] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, A. Prokudin and C. Turk, Phys. Rev. D **75**, 054032 (2007) [hep-ph/0701006]; M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, A. Prokudin and S. Melis, Nucl. Phys. Proc. Suppl. **191**, 98 (2009) [arXiv:0812.4366 [hep-ph]].

[50] F. Yuan and J. Zhou, Phys. Rev. Lett. **103**, 052001 (2009) [arXiv:0903.4680 [hep-ph]].

[51] Y. Koike, AIP Conf. Proc. **675**, 449 (2003) [hep-ph/0210396].

[52] A. Metz, Phys. Lett. B **549**, 139 (2002) [hep-ph/0209054].

[53] J. C. Collins and A. Metz, Phys. Rev. Lett. **93**, 252001 (2004) [hep-ph/0408249].

[54] F. Yuan, Phys. Rev. Lett. **100**, 032003 (2008) [arXiv:0709.3272 [hep-ph]]; Phys. Rev. D **77**, 074019 (2008) [arXiv:0801.3441 [hep-ph]].

[55] L. P. Gamberg, A. Mukherjee and P. J. Mulders, Phys. Rev. D **77**, 114026 (2008) [arXiv:0803.2632 [hep-ph]]; Phys. Rev. D **83**, 071503 (2011) [arXiv:1010.4556 [hep-ph]].

[56] S. Meissner and A. Metz, Phys. Rev. Lett. **102**, 172003 (2009) [arXiv:0812.3783 [hep-ph]].

[57] Y. Kanazawa and Y. Koike, Phys. Lett. B **478**, 121 (2000) [hep-ph/0001021]; Phys. Lett. B **490**, 99 (2000) [hep-ph/0007272].

[58] S. Meissner, Ph.D. thesis, University of Bochum, 2009.

[59] P. J. Mulders and R. D. Tangerman, Nucl. Phys. B **461**, 197 (1996) [Erratum-ibid. B **484**, 538 (1997)] [hep-ph/9510301].

[60] A. Bacchetta, M. Diehl, K. Goeke, A. Metz, P. J. Mulders and M. Schlegel, JHEP **0702**, 093 (2007) [hep-ph/0611265].